Hurricane Sandy in New York City

Lessons Learned



Some Lessons Learned . . .

- 1. Characterizing risk of major storms is a challenge
- 2. Hurricane Sandy grounded, reinforced, and motivated expansion of the NYC flexible adaptation approach
- 3. Federal regulations balkanized responses: 'The Tale of Two States and One City'
- 4. Federal, state, and municipal agencies need to coordinate on climate change scenarios
- 5. Having 'Science in place . . . Science in time' is critical

Hurricane Sandy as 'Tipping Point'



Hurricane Sandy Immediate Preparations







New York City issued mandatory evacuation of Zone A on October 28, 2012

Out-of-state utility crews brought in before the storm

MTA closed down operations, moved rolling stock, and boarded and placed sandbags at subway entrances to protect against flooding



SIRR, 2013

GISS Closes

Re: [giss-people-I] Sandy Columbia x









Ruedy, Reto A. (GISS-611.0)[TRINNOVIM, LLC] <reto.a.ruedy@nasa.gov>







GISS will be officially closed tomorrow; no decision has been made about Wednesday.

All windows are closed and printers were shut down. People currently at GISS are advised to leave early.

Thank you for your concern and cooperation,

Reto

Reto A. Ruedy Project Manager NASA Goddard Institute for Space Studies Trinnovim, LLC 2880 Broadway, Rm. 201A, New York, NY 10025 ph: 212-678-5541; email:reto.a.ruedy@nasa.gov WWUS71 KOKX 291617 NPW0KX URGENT - WEATHER MESSAGE NATIONAL WEATHER SERVICE NEW YORK NY 1217 PM EDT MON OCT 29 2012 ...HURRICANE FORCE WIND GUSTS THIS AFTERNOON AND EVENING...

A Few Hours Later . . .




Explosion at a substation on East 14th Street darkened most of Manhattan south of 34th Street.

Power remained on in Battery Park City, as Con-Ed did not shut service as network is on higher ground. More recent construction in Battery Park City provided buildings with generators and moved equipment to higher floors.

Despite Advanced Forecasts and Preparations . . .

- 44 deaths in NYC, 80% from drowning
- Major flooding 7 subway lines under East River, 3 tunnels closed
- 90,000 buildings in the inundation zone
- ~2 million without power
- Unforeseen impacts include gas shortages, hospital evacuations, and fires
- ~\$19 billion in damage in NYC




South Ferry Subway Station, Manhattan December 12, 2012


Source: D. Bader

SIRR, 2013

Hurricane Sandy


- Sandy followed an unusual storm track, turning sharply west just as it was reaching another peak of intensity
- Tropical-storm-force winds (39 to 73 mph) extended
 1,000 miles from end to end, making it more than
 three times the size of Hurricane Katrina
- Storms pressure was the lowest on record north of Cape Hatteras at 945mb
- **Storm timing** coincided almost exactly with astronomically high, high tide
- Storm surge combined with high tide created a storm tide of over 11 feet above NAVD88 at the Battery, highest water level on record (since 1856)







Comparing Flood Maps to Inundation Zone



FEMA 1983 100-year floodplain – ~200,000 New Yorkers

FEMA 2013 100-year floodplain – ~400,000 New Yorkers

FEMA 2013 100-year floodplain + NPCC2 90th percentile SLR projections for the 2050s

– ~800,000 New Yorkers


Outdated FEMA maps significantly underestimated the current risk level

Source: SIRR, 2013


Hurricane Sandy Links to Climate Change Science


Source: CCSR, 2013

- It is not possible to attribute any single extreme event such as Hurricane Sandy to climate change
- Sea level rise already occurring in the New York City area (~1.1 feet since 1900), in part related to climate change, increased extent and magnitude of coastal flooding during the storm
- Intensity of severe hurricanes in the northern hemisphere Atlantic Basin appears to be on rise and may increase in future***
- Melting sea-ice may be changing pattern of jet stream, making westward-turning storm tracks more likely***



North Atlantic SSTs and Hurricane Power Dissipation Index 1950-2005, Emanuel, 2007





Median Minimum Sea Ice Extent 1979-2009

***Areas of active research

Climate Change and a Global City 2001

Science in Place . . . Science in Time

The Potential

RESPONDING TO CLIMATE CHANGE
IN NEW YORK STATE
SYNTHESIS REPORT

ANNALS OF THE NEW YORK
ANNALS OF THE NEW YORK
ANNALS OF THE NEW YORK
1196

Climate Change
Adaptation in
New York City

Building a Risk Management Response

Potential future 1-in-100 yr flood zones for NYC using rapid ice melt model-based SLR projections

> 2020s 1-in-100 Year Flood Zone (w/ 9" Sea Level Rise) 2050s 1-in-100 Year Flood Zone

2080s 1-in-100 Year Flood Zone

Map Authors: K. Grady, A. Maroko, L. Patrick, W. Solecki, I For more information, contact: info@cunysusta

Interdependent Critical
Infrastructure Systems
and
Vulnerable Communities

South Ferry Subway Station



Beach erosion and boardwalk damage in the Rockaways







dan

Areas hard-hit by Sandy

Midland Beach Rockaways Coney Island Red Hook


New York City 2010 Report

Many impacts forecast well in advance


Sources: CCSR, 2013, MTA, 2012


Second New York City Panel on Climate Change

After Hurricane Sandy, Mayor Bloomberg re-convened the NPCC in January to provide updated climate risk information for the Special Initiative for Rebuilding and Resiliency (SIRR)



- The 2013 NPCC Climate Risk Information Report (CRI) provided new climate change projections and future coastal flood risk maps for New York City
- Both "A Stronger, More Resilient New York" and CRI reports released on June 11, 2013
- Full NPCC2 Report to be published in Fall 2014







NPCC2 Uncertainty and Risk Management

Projections


- Designed with NYC stakeholders to facilitate risk-based decision-making
- Based on results from 35 global climate models and 2 scenarios of future greenhouse gas emissions
- Direction of change associated with likelihoods as defined by the IPCC
- Qualitative projections provided for some extreme events, e.g., number of intense hurricanes, based on scientific literature and expert judgment





NPCC2 Climate Projection Methods Temperature and Precipitation


- Quantitative projections based on global climate model simulations (CMIP5)
 - 35 global climate models (GCMs)
 - 2 representative concentration pathways (RCP4.5, RCP8.5)
 - Timeslices: 2020s, 2050s, 2080s, 2100
 - 30-year average centered around decade
 - 1 ensemble member per GCM
- Downscaling
 - Delta method (Wilby et al., 2004)
 - Extreme events mean monthly changes applied to historical daily data
- Single gridbox method: Compare observed and model values for current climate; test changes in neighboring gridboxes; define as 100-mile radius; provide caveats upfront (Horton et al., 2001 JAMC)
- Projections provided for mean annual changes and extreme events



Data from IIASA

NPCC2 Climate Projection Methods Sea Level Change

- Innovative, 6-component approach used to develop regional sea level rise projections
 - Model-based components based on CMIP5 Data (24 GCMs and 2 RCPs)
 - Thermal expansion (global)
 - Changes in dynamic ocean height (local)
 - Components based on literature review and expert analysis
 - Ice mass loss from ice sheets and glaciers and ice caps (global)
 - Gravitational, rotational, and isostatic "fingerprinting" (local)
 - Vertical land movements (GIA) (local)
 - Land water storage (global)
- Projections are taken as the sum of the 6 sea level components
 - Provided for 2020s, 2050s, 2080s, 2100
 - 10-year average centered around decade
 - Changes are relative to the 2000 to 2004 base period




NPCC, 2013

Key Findings for Future Projected Changes

 Climate projections illustrate a broad-based acceleration of climate change in coming decades

- Show significant climate risks, especially heat waves, extreme precipitation events, and coastal flooding
- Valid for New York City and the metropolitan region
- By 2080s, projected changes include:
 - Annual temperature increase between 3.8°F to 10.3°F
 - Mean precipitation change between +5 and +15 percent
 - More likely than not increase in the number of the most intense hurricanes in the North Atlantic Basin.
 - Unknown how the total number of tropical cyclones and nor'easters will change in the North Atlantic Basin



NPCC, 2014 15

Mean Annual Changes – Sea Level Rise

Sea Level Rise Baseline (2000 – 2004)	Low-estimate (10 th percentile)	Middle range (25 th to 75 th percentile)	High-estimate (90 th percentile)
2020s	+ 2 in	+ 4 in to 8 in	+ 10 in
2050s	+ 8 in	+ 11 in to 21 in	+ 30 in
2080s	+ 13 in	+ 18 in to 39 in	+ 58 in
2100	+ 15 in	+ 22 in to 50 in	+ 75 in

Based on 24 GCMs and two Representative Concentration Pathways. Shown are the low-estimate (10th percentile), middle range (25th percentile to 75th percentile), and high-estimate (90th percentile).

Future Coastal Flood Heights and Recurrence Intervals at the Battery

2080s

	Current	Low- estimate (10 th percentile)	Middle range (25 th to 75 th percentile)	High-estimate (90 th percentile)
Annual chance of today's 100-year flood height	1 percent	1.7 percent	2.0 to 5.4 percent	12.7 percent
Today's 1-in-100 yr flood height to recur, on average	Once every 100 years	~ once every 60 years	~ once every 20 to 50 years	~ once every 8 years
Flood heights associated with 100-year flood	11.3 ft	12.4 feet	12.8 to 14.6 feet	16.1 feet


Flood heights are derived by adding the sea level rise projections for the corresponding percentiles to the baseline values. Baseline flood heights heights associated with the 100-year flood are based on the FEMA still water elevations, i.e., without wave height. Flood heights elevations are referenced to the NAVD88 datum.

Tale of Two States and One City


- Governor Cuomo of New York State established 2100
 Commissions immediately following Sandy; recognized climate change as increasing risk.
- Governor Christie of New Jersey focused on rebuilding 'as was;' does not recognize climate change as increasing risk
- Mayor Bloomberg, Leader of C40 Large Cities Climate Summit, created \$20 billion Resiliency and Rebuilding Plan, recognizes climate change as increasing risk
- US Congress required each entity to submit their own plans to receive appropriated funds; balkanized region

Risk Management Flexible Adaptation Pathways

Climate change adaptation as a risk management issue Flexible Adaptation Pathways as the response



Portfolio Approach to Resilience Action



Policy, social, engineering, and ecosystems interact to respond to changing climate and coastal hazards. Overlapping areas illustrate opportunities for adaptation and resilience strategies that combine components of each domain.

NYC SIRR Key Actions for Coastal Protection

Increase coastal edge elevations

 The City will increase the height of vulnerable coastal edges with beach nourishment and other measures over time




Minimize upland wave zones

 The City will work to provide significant attenuation of waves—that is, to knock down waves, or diminish their velocity—both off and onshore, before they reach neighborhoods

Protect against storm surge

 The City will use flood protection structures, such as floodwalls, levees, and local storm surge barriers built, where possible, to the 100-year flood elevation with an additional allowance for future sea level rise



Improve coastal design

 The City will study how natural areas and open space can be used to protect adjacent neighborhoods and maintain neighborhood quality of life, and will work to manage its own waterfront assets more effectively

Governance


Developing partnerships to improve permitting and study innovative coastal protections.

October, 2014 Update

5 of 37 coastal initiatives *complete*

35 of 37 coastal initiatives *in progress*

NOAA Coastal Map Tool



Source: NOAA; NPCC

NPCC Projections added to FEMA 2013 flood hazard areas in NOAA Sea Level Rise Map

22

Hurricane Sandy as Tipping Point

- Acceptable level of risk plummeted People are demanding greater protection from coastal storm risks
- Investment and action were mobilized by New York City
- NYC SIRR, US Congressional Appropriations, and Rebuild by Design program are creating a transformative new development trajectory
- Explicit recognition of increasing risks due to climate change in coastal areas now thoroughly accepted in public discourse
- Major effect on coastal storm protection planning not only in New York City, but in New York State, nationally, and even internationally.




Hartwig Kremer, May, 2014

Factors → Leadership; Federal/State/Municipal alignment; Media hub . . .

Looking forward . . .

Bloomberg target – 30% reduction of 2005 levels by 2050



Mayor de Blasio
has set
ambitious target
and timetable for
GHG reduction
and is
reconvening
NPCC, with
emphasis on
equity issues

City of New York, October 2014

"Global climate change is the challenge of our generation . . . New Yorkers will rise to the challenge. We will build on progress we have made to become more resilient to a changing climate and to mitigate the harmful greenhouse gas emissions that contribute to climate change. We are committing to reduce our emissions by 80 percent below 2005 levels by 2050, making us the largest city in the world to commit to this goal." Mayor De Blasio, Oct 2014


Temperature Change (2050s) and UCCRN Member Cities

City Size

Population of Metro Area


- Small (<500,000)
- Intermediate (500,000-1,000,000)
- Big (1,000,000-5,000,000)
- Large (5,000,000-10,000,000)
- Mega (>10,000,000)

Temperature Change (Degrees C)



4.5 - 5.25

>5.25



UCCRN Member Cities

			Australia/Ossania	
<u>Africa</u>	<u>Asi</u>	<u>a</u>	Australia/Oceania	
Abuja Cape Town Dakar Durban Harare Johannesburg Kampala	Bangkok Beijing Chennai Delhi Dhaka Eskisehir Hong Kong	Sargodha Seoul Shanghai Tokyo Ulaanbaatar	Gold Coast Melbourne Parkville Sydney Townsville Wellington Wembley	
Nairobi	Jaipur			
Rabat Setif Sfax Lagos	Kathmandı Kyoto Nagoya Ningbo	I		
*colors represent mean annual temperature				

*colors represent mean annual temperature change for a mid-range scenario (RCP4.5) from CMIP5 models (2040-2069 average minus 1971-2000 average).

Euro	<u>pe</u>
Aalborg	Leipzig
Athens	London
Barcelona	Luxembourg
Berlin	Naples
Bonn	Newcastle upon Tyn
Bristol	Oxford
Brussels	Paris
Copenhagen	Peterborough
Enschede	Planken
Exeter	Potsdam
Freiburg	Rome
Geneva	Stockholm
Glasgow	Stuttgart
Groningen	Tallinn
Helsinki	Trieste
Istanbul	Venice

Kokkola

North America			
Amherst	Mexico City		
Atlanta	Montreal		
Aurora	Mountain View		
Baton Rouge	New Haven		
Boston	New Orleans		
Boulder	New York		
Cambridge	Norfolk		
College Park	North Little Rock		
College Station	ı Nyack		
East Lansing	Ottawa		
Englewood	Reno		
Eugene	Sacramento		
Guelph	Saint Catherines		
Hauppauge	San Diego		
Idaho Falls	Seattle		
Kingston	Toronto		
Los Altos	Tucson		
Los Angeles	Washington DC		
Martinez	Yardley		

South America Brasilia Buenos Aires Concón Curitiba Lima Montevideo Rio de Janeiro Santa Cruz Santiago Sao Paulo

(35 CMIP5 models)

Vienna

References and Links

- Consortium for Climate Risk in the Urban Northeast (<u>www.ccrun.org</u>)
- NYSERDA ClimAID (<u>www.nyserda.ny.gov/climaid</u>)
- New York City Panel on Climate Change report available online at (<u>www.nyas.org</u>)
- Urban Climate Change Research Network (<u>www.uccrn.org</u>)









